

CHAPTER 3

Nuclear Models

Lecture Notes For

PHYS 415

Introduction to Nuclear and Particle Physics

To Accompany the Text

Introduction to Nuclear and Particle Physics, 2nd Ed.

A. Das and T. Ferbel

World Scientific

General Remarks

- The nuclear force has proved elusive.
 - Scattering experiments gave many clues, but, at the time, no fundamental theory of the nuclear force existed.
 - Even when the fundamental theory was developed (QCD), due to the strong coupling nature of the force, first principle calculations were impossible.
 - Lately, various techniques have been developed, such as lattice QCD, but, currently, they too are limited in what they are able to predict.
- Nuclear models were developed as a result.
 - Phenomenological basis.
 - Limited range of validity.

Liquid Drop Model


- Nuclear densities are almost independent of nucleon number:

$$R = r_0 A^{1/3} \Rightarrow \rho = \frac{M}{V} \approx \frac{m_N A}{\frac{4}{3} \pi R^3} = \frac{m_N A}{\frac{4}{3} \pi r_0^3 A} = \frac{m_N}{\frac{4}{3} \pi r_0^3} = \text{constant}$$

- This suggests the nucleus as an incompressible fluid: **liquid drop**.
- Adding more nucleons, increases the size, but not the density.
 - Only nearest neighbor interactions are important.
 - Nuclear force *saturates*.
 - Consistent with $B/A \approx \text{constant}$.

Surface Tension

- Nucleons at the surface feel forces only from interior nucleons.
 - The force is unbalanced at the surface.
 - Since the force is attractive, there is a net inward attraction of the surface: *surface tension*.
 - Surface nucleons are less tightly bound.




Binding Energy, Including Surface Effect

- Less tightly bound surface nucleons imply a positive correction to the binding energy:

$$\text{B.E.} = -a_1 A + a_2 A^{2/3}$$

Volume energy



Surface correction

- Surface effect is more important for lighter nuclei.
 - Higher surface-to-volume ratio
 - Therefore, light nuclei are less tightly bound

Coulomb Forces

- The coulomb repulsion among protons decreases the binding energy per nucleon:

$$\text{B.E.} = -a_1A + a_2A^{2/3} + a_3 \frac{Z^2}{A^{1/3}}$$

since $R \propto A^{1/3}$

Bethe-Weizsäcker Semi-Empirical Mass Formula

- We still need to account for other observations:
 - Light nuclei with $N = Z$ are more abundant (stable)
 - Even-even nuclei are more abundant (stable).

$$\text{B.E.} = -a_1 A + a_2 A^{2/3} + a_3 \frac{Z^2}{A^{1/3}} + a_4 \frac{(N - Z)^2}{A} \pm a_5 A^{-3/4}$$

$$\Rightarrow M(A, Z) = (A - Z)m_n + Zm_p$$

$$- \frac{a_1}{c^2} A + \frac{a_2}{c^2} A^{2/3} + \frac{a_3}{c^2} \frac{Z^2}{A^{1/3}} + \frac{a_4}{c^2} \frac{(A - 2Z)^2}{A} \pm \frac{a_5}{c^2} A^{-3/4}$$

where the last term is

$$\left\{ \begin{array}{l} + \text{ For odd-odd nuclei} \\ - \text{ For even-even nuclei} \\ 0 \text{ For odd } A \text{ nuclei} \end{array} \right.$$

and, from an empirical fit


$a_1 \approx 15.6 \text{ MeV}$	$a_2 \approx 16.8 \text{ MeV}$	$a_3 \approx 0.72 \text{ MeV}$
$a_4 \approx 23.3 \text{ MeV}$	$a_5 \approx 34 \text{ MeV}$	

Fermi-Gas Model

- Quantum-mechanical description where nucleons are considered a gas of fermions confined to a spherically symmetric potential well.
- Boundary conditions imply energy levels are discrete.
- Depth and range of well are fit to data.
- Only two identical fermions (of opposite spin projection) can occupy the same energy level.
 - Protons and neutrons may be considered distinguishable and so each level can contain four nucleons.
 - Protons experience the Coulomb force and so the potentials are slightly different for protons vs. neutrons.

Fermi Levels and Fermi Energy

- Protons occupy shallower well since heavy nuclei are neutron rich. Otherwise, neutrons in upper levels could undergo β^- decay to become lower energy protons.



Fermi Energy

- Nonrelativistically, $E_F = p_F^2 / 2m$
- Phase space volume = $\int d^3r d^3p$:

$$V_{TOT} = V \times V_F = \frac{4\pi}{3} r_0^3 A \times \frac{4\pi}{3} p_F^3 = \left(\frac{4\pi}{3}\right)^2 (r_0 p_F)^3 A$$

- Heisenberg: $\Delta x \Delta p_x \geq \frac{\hbar}{2} \Rightarrow V_{\text{state}} = (2\pi\hbar)^3 = h^3$
- Number of fermions up to E_F :

2 spin states

$$n_F = 2 \frac{V_{TOT}}{V_{\text{state}}} = \frac{4}{9\pi} A \left(\frac{r_0 p_F}{\hbar} \right)^3$$

Fermi Energy, cont'd.

- Assuming $N = Z = A / 2$:

$$N = Z = \frac{A}{2} = \frac{4}{9\pi} A \left(\frac{r_0 p_F}{\hbar} \right)^3 \Rightarrow p_F = \frac{\hbar}{r_0} \left(\frac{9\pi}{8} \right)^{1/3}, \text{ indep. of } A$$

$$\Rightarrow E_F = \frac{1}{2m} \left(\frac{\hbar}{r_0} \right)^2 \left(\frac{9\pi}{8} \right)^{2/3} \approx \frac{2.32}{2mc^2} \left(\frac{\hbar c}{r_0} \right)^2 \approx 33 \text{ MeV}$$

- Taking B.E. of last nucleon as ~ 8 MeV:

$$V_0 = E_F + B \approx 40 \text{ MeV}$$

Shell Model

- Nucleons within nuclei, like electrons within atoms, can be well described by a **shell model**.
- Four nucleons (two protons and two neutrons) can occupy each orbital.
- For atoms, this is natural, since the nucleus provides a central force. For nuclei there is no central force!
- The Pauli principle still forbids nucleons from occupying the same states and this gives rise to an effective **mean field potential** in which the nucleons move.

Review of Atomic Shell Model

- Electrons occupy states defined by four quantum numbers: n, ℓ, m_ℓ, m_s
 - $n = 1, 2, 3, \dots$ = principal quantum #
 - $\ell = 0, 1, 2, \dots, n-1$ = orbital quantum #
 - $m_\ell = -\ell, -\ell + 1, \dots, 0, 1, \ell-1, \ell$ = magnetic quantum #
 - $m_s = \pm 1/2$
- Considering only the nuclear Coulomb potential, which is rotationally symmetric, all states of a given n are *degenerate*: # of states = $2n^2$.

Spin-Orbit Interaction

- In atoms, the spin of an electron couples to the nuclear orbital motion (as seen in the electron rest frame), giving rise to small splittings: *fine structure*.
- In nuclei, there is also a spin-orbit interaction and this affects nuclear structure significantly.

Closed-Shell Atoms

- In atoms there is a strong pairing effect which effectively minimizes the angular momentum of the ground state.
 - $\sum m_s = 0$ and $\sum m_l = 0$
 - $\vec{L} = 0 = \vec{S}$ and $\vec{J} = \vec{L} + \vec{S} = 0$
- Closed shell elements are very stable.
 - Chemically inert
 - Large ionization energies

Magic Numbers

- Atomic shell closures occur at certain *magic numbers*: $Z = 2, 10, 18, 36, 54$
- Nuclear shell closures also occur at well-defined magic numbers:
 - $N = 2, 8, 20, 28, 50, 82, 126$
 - $Z = 2, 8, 20, 28, 50, 82$
- Nuclei where both protons and neutrons have closed shell are called **doubly magic** and have even greater stability.

Other Evidence for Nuclear Shells

- Closed neutron shell nuclei have more *isotones* and closed proton shell nuclei have more *isotopes* than neighboring nuclei.
- Neutron capture cross sections are relatively small for closed-shell nuclei.
- Proton and neutron knockout experiments can probe nucleons in individual orbits.
 - Can measure binding energies this way.
 - Can deduce the momentum distribution (in the context of a reaction model). The Fourier Transform of this gives the shape of the orbital.

Schrödinger Equation and Solutions

- For a central potential:

$$\left(-\frac{\hbar^2}{2m} \vec{\nabla}^2 + V(r) \right) \psi(\vec{r}) = E\psi(\vec{r})$$

- Since the potential is spherically symmetric:

$$\left[H, \vec{L}^2 \right] = 0 \Rightarrow \text{sol'ns. are simultaneous eigenstates of } H \text{ and } \vec{L}^2$$

- Solutions can be written as:

$$\psi_{nlm_\ell}(\vec{r}) = \frac{u_{nl}(r)}{r} Y_{\ell m_\ell}(\theta, \phi)$$

Radial and Angular Equations

- Angular equations:

$$\vec{L}^2 Y_{\ell m_\ell}(\theta, \phi) = \ell(\ell + 1)\hbar^2 Y_{\ell m_\ell}(\theta, \phi)$$

$$L_z Y_{\ell m_\ell}(\theta, \phi) = m_\ell \hbar Y_{\ell m_\ell}(\theta, \phi)$$

- Radial equation:

$$\left[\frac{d^2}{dr^2} + \frac{2m}{\hbar^2} \left(E_{n\ell} - V(r) - \underbrace{\frac{\hbar^2 \ell(\ell + 1)}{2mr^2}} \right) \right] u_{n\ell}(r) = 0$$

Centrifugal barrier



Boundary conditions: $u_{n\ell}(r) \xrightarrow{r \rightarrow 0, \infty} 0$

Parity

- The parity operation is defined by: $P\psi(\vec{r}) = \psi(-\vec{r})$

- This is equivalent to:
$$\begin{cases} r \rightarrow r \\ \theta \rightarrow \pi - \theta \\ \phi \rightarrow \pi + \phi \end{cases}$$

- For spherically symmetric potentials, states have definite parity, related to the value of ℓ :

$$\text{parity} = (-1)^\ell$$

$$\Rightarrow \text{parity} = \quad + \text{ for } \ell \text{ even}$$

$$\quad - \text{ for } \ell \text{ odd}$$

Infinite Square Well

- We can gain some intuition by solving this equation for simple potentials, even though they might not be entirely reasonable.
- For the ∞ square well:

$$V(r) = \begin{cases} \infty & r \geq R \\ 0 & \text{otherwise} \end{cases}$$

- We require that the solutions be regular at the origin, so we get the *spherical Bessel functions*:

$$u_{n\ell}(r) = j_{\ell}(k_{n\ell}r) \quad \text{where} \quad k_{n\ell} = \sqrt{\frac{2mE_{n\ell}}{\hbar^2}}$$

Energies are Quantized

- Since the potential is infinite at $r = R$, we must have:

$$u_{n\ell}(R) = j_{\ell}(k_{n\ell}R) = 0, \quad \ell = 0, 1, 2, 3, \dots$$

and $n = 1, 2, 3, \dots$ for any ℓ

- The energies are quantized and depend on n **and** ℓ . Each energy level can contain $2(2\ell + 1)$ protons or neutrons.
- For $n = 1$, we get shell closures at:
2, 2+6=8, 8+10=18, 18+14=32, 32+18=50, ...
(observed: 2, 8, 20, 28, 50, 82, ...)
- Some, but not all, magic numbers are reproduced.

Harmonic Oscillator

- The three dimensional harmonic oscillator:

$$V(r) = \frac{1}{2} m \omega^2 r^2$$

- Solutions are related to the *associated Laguerre polynomials* with additional Gaussian factor (dominant dependence for large r).
- We also get bound states with discrete energies:

$E_{nl} = \hbar\omega \left(2n + \ell - \frac{1}{2} \right)$ <p>and $\ell = 0, 1, 2, \dots$, for any $n = 1, 2, \dots$</p>	$\Lambda \equiv 2n + \ell - 2$ \longrightarrow	$E_{nl} = \hbar\omega \left(\Lambda + \frac{3}{2} \right)$ <p>$\Lambda = 0, 1, 2, \dots$</p>
---	---	--

Shell Closures

- Levels with different (n, ℓ) , leading to the same Λ , will be degenerate. The degeneracy is:

$$n_{\Lambda} = (\Lambda+1)(\Lambda+2)$$

- Therefore, shell closures occur for proton or neutron numbers of 2, 8, 20, 40, 70, ...

(observed: 2, 8, 20, 28, 50, 82, ...)

- Again, some, but not all, magic numbers are reproduced.
- So introduce ...

Spin-Orbit Potential

- Maria Goeppert Mayer and Hans Jensen (1949) suggested adding a spin-orbit interaction, in analogy with atoms:

$$V_{\text{TOT}} = V(r) - f(r)\vec{L} \cdot \vec{S}$$

- Similar to atoms, except
 - Presence of $f(r)$ function
 - $j = \ell + 1/2$ state has lower energy than $j = \ell - 1/2$

Splitting due to Spin-Orbit Interaction

- We can write $\vec{L} \cdot \vec{S} = \frac{1}{2}(\vec{J}^2 - \vec{L}^2 - \vec{S}^2)$

$$\Rightarrow \Delta = \Delta E_{nl} \left(j = \ell - \frac{1}{2} \right) - \Delta E_{nl} \left(j = \ell + \frac{1}{2} \right)$$

$$= \hbar^2 \left(\ell + \frac{1}{2} \right) \int d^3r |\psi_{nl}(\vec{r})|^2 f(r)$$

- Splitting is larger for higher ℓ values, allowing level crossing.
- For appropriate $f(r)$, we can reproduce all the magic numbers.


Spectroscopic Notation

- As for atoms, we use the spectroscopic notation to label states:

$$nL_j$$

- The L value is labeled by S, P, D, F, G, \dots , for $\ell = 0, 1, 2, 3, 4, \dots$
- The multiplicity is $2j+1$

Level Scheme with Spin-Orbit Term



From: <http://hyperphysics.phy-astr.gsu.edu/hbase/nuclear/shell.html>

Predictions of the Shell Model

- Spin-parity assignments of ground states of many odd- A nuclei predicted.
 - Neutrons and protons pair up with opposite spins, so that the spin of the last nucleon determines the spin of the nucleus.
 - Even-even nuclei, consequently, have zero spin, in agreement with observations.
- Allowing pairing between all valence nucleons can fix up agreement with remaining odd- A nuclei.

Example of Spin-Parity Prediction

- Consider $^{13}\text{C}^6$.
- Six protons and six neutrons are completely paired off.
- The last neutron will be in the shell: $1P_{1/2}$.
- So we expect $(1/2)^-$ (i.e. $j = 1/2$ and negative parity, since ℓ is odd).
- This is in agreement with observation.

Magnetic Moments of Nuclei

- We expect the nuclear magnetic moment to be determined by the moment of the unpaired nucleon(s).
- Each unpaired nucleon contributes:
 - Intrinsic (spin): $\mu_p = 2.79 \mu_N$ and $\mu_n = -1.91 \mu_N$
 - Orbital: protons only; neutrons are uncharged.
- Example: deuteron, assuming proton and neutron are in $(1S)_{1/2}$ states:
 - Prediction: $\mu_d = 2.79 \mu_N - 1.91 \mu_N = 0.88 \mu_N$
 - Observation: $0.86 \mu_N$
- Example: ${}^3\text{He}^2$. Here, the magnetic moment is expected to be due to the unpaired neutron and this is roughly correct. This fact, has been exploited to use ${}^3\text{He}^2$ targets to deduce neutron properties.

Collective Model

- The single-particle shell model does not describe certain features of heavy nuclei, namely magnetic dipole moments and electric quadrupole moments.
 - Electric quadrupole moments arise from non-spherical charge distributions, which cannot be explained by a purely central potential.
- Including many-body physics where nucleons interact with one another can remedy the problem.
- Historically, various collective models were introduced which provided simpler, intuitive descriptions of heavy nuclei.

Deformed Rotators and Vibrators

- Aage Bohr, Ben Mottelson and James Rainwater proposed a collective model to explain observed moments.
- Tried to reconcile liquid drop and shell model.
 - Hard core of nucleons in filled shell-model states.
 - Surface motion (rotation) of valence nucleons.
 - Latter gives rise to nonspherical shape and rotational and vibrational energy spectra.

Ellipsoidal Nucleus

- Define surface of nucleus as:

$$ax^2 + by^2 + \frac{z^2}{ab} = R^2$$

- Mean potential given by:

$$V(x, y, z) = \begin{cases} 0 & \text{for } ax^2 + by^2 + \frac{z^2}{ab} \leq R^2 \\ \infty & \text{otherwise} \end{cases}$$

- Deformations into an ellipsoidal shape can also be induced through particle bombardment of heavy nuclei. This will be useful for discussing nuclear fission later.

Rotational and Vibrational Levels

- For rotations, define the Hamiltonian:

$$H = \frac{\vec{L}^2}{2I}, \quad \text{with eigenvalues} = \frac{\ell(\ell + 1)}{2I} \hbar^2$$

- For rotations perpendicular to symmetry axis, only expect even ℓ .
- Photon quadrupole transitions ($\Delta\ell = 2$) have been observed, corresponding to transitions between rotational levels.

Superdeformed Nuclei

- Heavy ion collision experiments have produced superdeformed nuclei with very large angular momentum quantum numbers.
- These deformed nuclei reach more spherical shapes by emitting a series of quadrupole γ -rays, each of order 50 keV.
- The photon energies remain essentially fixed. This is in conflict with the collective model, since the decreasing moment of inertia during “spin-down” should give rise to nonuniformly spaced photon energies.
- Emissions from different nuclei are also nearly identical. This too is in conflict with current models since effects of nucleon pairing should produce varying level spacings.